
New Correction Method



Function approximation method using ratei(3) and TIS

Let's take a relatively straightforward idea...

$$F_{i=1...3}(TIS) \cdot RateI(3) + A_{0}$$


$$F_{i}(TIS) \equiv A_{2i} \cdot TIS + A_{2i+1}$$
[range]
$$F_{i} \equiv \begin{cases} F_{1} & TIS \leq 0.5 \\ F_{2} & 0.5 < TIS \leq 2.0 \\ F_{3} & 2.0 < TIS \end{cases}$$

Parameter size:

7(parameters) * 912(PMT) = 6384


Cf. matrix : 570000

New Correction Method -result



New Correction Method -result

Categorization of miss correction patterns.



Is there any other miss correction?

Any inclination to fail?

Eye scan
And categorize them.


1: fail at the spill end.

2: success to correct.

New Correction Method – result cont...

Let's take a look at more global things...

The sigma/mean ratio over the up-stream PMTs




X-axis: PMT Number (up stream)

NOTE: Mean is normalized to 1000. rsmon.raw00.50007.hbook

New Correction Method – result cont...


Can we apply this function (and its parameters) to another run?



Rsmon.raw00.50007.hbook

Rsmon.raw00.50061.hbook

Correction Power of Jim's Matrix Method




Can new function method have enough correction power compare to the Jim's matrix method?

Do matrix based correction using the same data sets, and Compare!

There is a tendency to rectify them to the miss direction at the both edges of spill structure.

rsmon.raw00.50061.hbook

Comparison between New correction Method and Jim's Matrix Method



New correction method has the almost same correction effect as Matrix method.